

Propel/Miralax Bowel Prep

You will need to drink a laxative solution to clean your colon. Please purchase the following items over the counter:

- General Genera
- Dulcolax laxative tablets (not suppository or stool softener). You will need 3 tablets for the prep.
- Miralax 238 grams (8.3 ounces) powder or generic polyethylene glycol 3350 (you can find this in the laxative section).
- One Reglan Tablet (generic Metoclopramide) 10 mg (prescription only)

One day before your colonoscopy:

NO SOLID FOOD - Clear liquids only for breakfast, lunch & dinner. Please make an effort to drink clear liquids throughout the day.

- 1. Start clear liquids as soon as your get up: Clear liquid list provided below. No Solid Foods.
- 2. In the morning, in a pitcher, mix the 8.3-ounces of Miralax with the 64-ounces of Propel. Stir/shake the contents until entire contents of Miralax are completely dissolved. Chill if desired.
- 3. At 1:00 pm: Take three Dulcolax laxative tablets with water by mouth
- 4. At 2:00 pm: Take the Reglan tablet
- 5. At 3:00 pm: Drink ½ of the Propel/Miralax solution. Drink 8-ounce glass every 15 minutes. You may drink it directly through a straw.
- 6. At 9:00 pm: Drink the other ½ of the Propel/Miralax solution. Drink 8-ounce glass every 15 minutes until finished with all of the solution.
- 7. Clear liquids allowed until midnight and then nothing to eat or drink.

Day of Procedure:

- Nothing to drink
- Have someone with you to drive you home

Clear Liquid Diet List - Do not eat or drink anything colored red or purple

- Beverages: Gatorade, Kool-Aid, Strained fruit juices without pulp apple, white grape, orange, lemonade, Water, tea or coffee (no milk, non-dairy creamer, or lemon)
- Broths: Chicken/beef bouillon or vegetable broth
- Desserts: Hard candies, Jell-O (lemon, lime or orange; no fruit or toppings) Popsicle (no sherbet or fruit bars)

÷¢